

Figure 1. Recommended immunization schedule for adults aged 19 years or older by age group, United States, 2017

| Vaccine                        | 19–21 years   | 22–26 years | 27–59 years | 60–64 years | ≥ 65 years |
|--------------------------------|---|-------------|-------------|-------------|------------|
| Influenza <sup>1</sup>         | 1 dose annually   |             |             |             |            |
| Td/Tdap <sup>2</sup>           | Substitute Tdap for Td once, then Td booster every 10 yrs |             |             |             |            |
| MMR <sup>3</sup>               | 1 or 2 doses depending on indication                      |             |             |             |            |
| VAR <sup>4</sup>               | 2 doses   |             |             |             |            |
| HZV <sup>5</sup>               |   |             |             | 1 dose      |            |
| HPV–Female <sup>6</sup>        | 3 doses   |             |             |             |            |
| HPV–Male <sup>6</sup>          | 3 doses   |             |             |             |            |
| PCV13 <sup>7</sup>             |   |             |             |             | 1 dose     |
| PPSV23 <sup>7</sup>            | 1 or 2 doses depending on indication                      |             |             |             | 1 dose     |
| HepA <sup>8</sup>              | 2 or 3 doses depending on vaccine                         |             |             |             |            |
| HepB <sup>9</sup>              | 3 doses   |             |             |             |            |
| MenACWY or MPSV4 <sup>10</sup> | 1 or more doses depending on indication                   |             |             |             |            |
| MenB <sup>10</sup>             | 2 or 3 doses depending on vaccine                         |             |             |             |            |
| Hib <sup>11</sup>              | 1 or 3 doses depending on indication                      |             |             |             |            |



Recommended for adults who meet the age requirement, lack documentation of vaccination, or lack evidence of past infection



Recommended for adults with additional medical conditions or other indications



No recommendation